



# FERTILIZATION GUIDE FOR BEAN

## FERTILIZATION GUIDE FOR BEAN



### BEAN FERTILIZATION

The numbers on the fertilizer bags indicate by weight what percentages of each of the three most important nutrients - nitrogen (N), phosphorus (P) and potassium (K) - are present in the mixture. Although the percentages may vary, the order is always the same: N, P and K.

Nitrogen promotes healthy green leaves and stems, and you don't need much of it for beans. If you have too much, the plant will spend more time making leaves and less time making beans. Phosphorus promotes strong roots and potassium conditions the whole plant, helping it to bear fruit and resist disease.



# FERTILIZATION GUIDE FOR BEAN



## DRIP IRRIGATION FERTILIZATION PROGRAM

### Base Fertilization

Pulcu Leo Bio	40 - 50 Kg
Pulcu Excellent 14-12-14+13S	40 - 50 Kg



### Drip Irrigation

After Planting	Per Acre
2P AminoMax 45	100 cc
Nano Elite 30-10-10+Te	2 Kg
Pulcu Expert 10-52-10+Te	2 Kg
Following the Development	Per Acre
2P Nano Root	500 cc
Pulcu Expert 20-20-20+Te	2.5 Kg
During the fruit growing period	Per Acre
2P AminoMax 45	100 cc
Pulcu Expert 16-8-24+Te	2.5 Kg
During the Harvest Period	Per Acre
2P Nano Tek	100 cc
Nano Elite 15-5-40+Te	2.5 Kg

Pulcu Expert 10-52-10+TE should be used if there is a weakening (decrease) in the flowers during the production process.

# FERTILIZATION GUIDE FOR BEAN



## FOLIAR FERTILIZATION PROGRAM

To Promote Flower Formation	Per Acre
2P Nano Root	200 cc
2P Zinc EDTA 15%	80 gr
Pulcu Expert 10-52-10+Te	200 gr
If Vegetative Part (Green Part) Is Weak	Per Acre
2P Cytokinin	50 cc
2P Gold Combi	60 gr
Pulcu Expert 20-20-20+Te	200 gr
For Fruit Development	Per Acre
2P BorMax	150 cc
Nano Elite 17-7-21+6,5CaO+2,5MgO+TE	200 gr
Against Fruit Softening and Flower Nose Rot	Per Acre
2P Cal - L	250 cc
For Color, Quality and Size of Fruit	Per Acre
2P Cytokinin	100 cc
Pulcu Expert 7-6-40+TE	250 gr
2P Nano Tek	250 cc

For the vegetative and generative development of the plant, immediately after planting, 100 lt of water / 100 cc 2P CYTOKIN should be applied 3 times with an interval of 5 days.





**MANUFACTURING**  
**PREMIUM QUALITY**  
**PESTICIDES & FERTILIZERS**